



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/00	A2	(11) International Publication Number: WO 99/44581 (43) International Publication Date: 10 September 1999 (10.09.99)
(21) International Application Number: PCT/US99/04449 (22) International Filing Date: 1 March 1999 (01.03.99) (30) Priority Data: 60/076,770 4 March 1998 (04.03.98) US (71) Applicant (for all designated States except US): OR- THO-McNEIL PHARMACEUTICAL, INC. [US/US]; U.S. Route #202, Raritan, NJ 08869-0602 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): THAKUR, Madhav, S. [US/US]; 112 Mendham Drive, North Wales, PA 19477 (US). KOTWAL, Pramod, M. [US/US]; 145 Cambronne Circle, Blue Bell, PA 19422 (US). GIBBS, Irwin, S. [US/US]; 144 Roy Lane, Huntington Valley, PA 19006 (US). (74) Agents: CIAMPORCERO, Audley, A., Jr. et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08903-7003 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PHARMACEUTICAL COMPOSITION OF TOPIRAMATE (57) Abstract The invention is directed to a pharmaceutical composition of topiramate, an anticonvulsant which is useful for treating epilepsy. More specifically, the present invention provides a solid dosage formulation of topiramate intended primarily for use by pediatric patients, of for patients who have difficulty swallowing tablets. Processes for preparing the pharmaceutical composition are also described.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title of the Invention

Pharmaceutical Composition of Topiramate

5 Cross-Reference to Related Application

This application claims priority from United States provisional application Serial No. 60/076,770, filed March 4, 1998, the contents of which are hereby incorporated by reference.

10

Field of the Invention

The present invention provides a solid dosage formulation of topiramate and process for producing the solid dosage formulation. More particularly, the solid dosage formulation comprises core particles which are coated with a taste mask coating to provide coated particles which can be sprinkled onto food to ease administration to patients who have difficulty swallowing tablets or capsules, e.g., pediatric patients.

20 Background of the Invention

The pharmaceutical industry employs a variety of dosage formulations for orally administering medicinal agents to patients. Typical formulations for oral administration include liquid solutions, emulsions, or suspensions, as well as solid forms such as capsules or tablets (as used herein, the term "tablet" means any shaped and compressed solid dosage form, including caplets). Since these conventional solid dosage formulations are usually intended for adults who can easily swallow large tablets whole, the often disagreeable taste of the active ingredient need not be taken into account in formulating the medicine, except for the provision of means to prevent the taste from being apparent during the short time that the medicine is in the mouth. Such means may include the provision of an appropriate coating on the tablet, the use of a capsule form (the gelatin outer

shell of the capsule keeps the active ingredient inside until the capsule has been swallowed), or simply firmly compressing a tablet so that it will not begin to disintegrate during the short time that it is intended to be in the mouth.

5 Children, older persons, and many other persons have difficulty swallowing whole tablets and even capsules. Therefore, it is often desirable to provide the medicine either in liquid form or in a chewable solid form or an alternative solid form, e.g., small particles which can be sprinkled onto soft food and swallowed
10 intact with the food, in addition to the tablet or capsule intended to be swallowed whole. Even where the medicine can be formulated as a liquid, it is desirable to provide a chewable solid form or an alternative solid form such as microspheres which can be sprinkled onto soft food (e.g., baby food) because it is often more convenient and easier to administer.

15 A major requirement of any such solid form is that it must be palatable, since an unpalatable formulation greatly increases the risk of a patient neglecting to take a medication. A further requirement of any solid dosage form is that it must be bioavailable; that is, once the formulation reaches the stomach, the individual particles should release the active ingredient rapidly and completely to
20 ensure that substantially all of the active ingredient is absorbed. In cases where the active ingredient is particularly unpalatable and somewhat unstable, it may be difficult, if not impossible, to identify a solid form that fulfills both of these requirements (i.e., palatable and bioavailable).

25 A number of references are known which describe pharmaceutical compositions of unpalatable medicinal agents which are coated with a taste masking coating in order to hide the unpleasant taste. Julian et al., in U.S. Patent No. 4,851,266, describe chewable medicament tablets made by coating granules of a medicament (especially, acetyl p-aminophenol) with a blend of cellulose
30 acetate or cellulose acetate butyrate and polyvinyl pyrrolidone (also known as "PVP" and referred to hereinafter by its United States Pharmacopeia (USP) name as "povidone"). Mehta, in U.S. Patent No. 5,084,278, discloses a pharmaceutical

composition comprised of a pharmaceutical core of an active dose of a compound and a microencapsulating polymer which coats the pharmaceutical core and is capable of taste-masking the active compound. Bhardway, et al., in U.S. Patent No. 5,578,316, describe medicament cores coated with methacrylate ester
5 copolymers which mask the bitter and unpleasant taste of the medicament.

A variety of chlorosulfate and sulfamate esters of 2,3:4,5-bis-Q-(1-methylethylidene)- β -D-fructopyranose, and their anticonvulsant activity in mammals, and thus their utility in treating diseases such as epilepsy and
10 glaucoma, are described in U.S. Patent No. 4,513,006. More specifically, the compound 2,3:4,5-bis-Q-(1-methylethylidene)- β -D-fructopyranose sulfamate, hereinafter referred to as "topiramate", is presently available for marketing as a tablet product in strengths of 25, 50, 100, 200, 300 and 400 mg as adjunctive therapy for the treatment of adults with partial onset seizures (TOPAMAX®
15 (topiramate) tablets). Topiramate can be prepared following the processes disclosed in U.S. Patent Nos. 4,513,006 and 5,387,700, and preferably, by the process described in Examples 1 to 3 of U.S. Patent No. 5,387,700. Difficulty in identifying a chewable solid form of topiramate has ensued due to the extremely bitter taste of topiramate and problems associated with stability of the active agent,
20 especially upon exposure to moisture and heat which are known to cause degradation of topiramate. Degradation of topiramate is readily detected by changes in physical appearance, i.e., discoloration to brown or black, and by the formation of sulfate ions which can be readily detected by standard techniques know to those of ordinary skill in the art (e.g., HPLC).

25

Accordingly, it is an object of the invention to provide a stable solid formulation of topiramate for use in children and other patients who have difficulty swallowing conventional solid forms (e.g., tablets, capsules) which is both palatable and bioavailable. It is a further object of the invention to provide a
30 palatable solid formulation of topiramate that can be sprinkled onto soft food prior to consumption (i.e., a "sprinkle formulation") and which provides immediate release of the active ingredient in the stomach.

Summary of the Invention

The present invention is directed to a process for forming a pharmaceutical
5 composition comprising:

- (a) preparing core particles comprising an active agent of topiramate;
- (b) drying the core particles from step (a) to form dried core particles;
- (c) coating the dried core particles from step (b) with a taste masking
mixture to form coated particles; and
- 10 (d) drying the coated particles from step (c) to form the pharmaceutical
composition wherein the amount of taste masking mixture ranges from about 7%
by weight to about 15% by weight of the pharmaceutical composition, preferably,
about 9 to about 13%, and most preferably, about 11% by weight of the
pharmaceutical composition.

15

In another aspect of the invention is a pharmaceutical composition
comprising:

- (a) core particles containing an active agent of topiramate, wherein the core
particles have an initial particle size between about 0.100 mm and 2.5 mm; and
- 20 (b) a taste mask coating, wherein the taste mask coating comprises
between about 7% by weight and about 15% by weight of the pharmaceutical
composition, preferably, about 9 to about 13%, and most preferably, about 11% by
weight of the pharmaceutical composition and wherein the coated particles of the
pharmaceutical composition have a final particle size of about 0.100 mm to about
25 2.5 mm.

In one embodiment of the invention, the core particles comprise the active
agent of topiramate and at least one excipient; preferably, the core particles
comprise the active agent of topiramate, a binder and a diluent; more preferably,
30 the core particles comprise the active agent of topiramate, a binder and sugar
spheres.

In a class of the invention is a pharmaceutical composition comprising about 85 to about 93% by weight core beads, and about 7 to about 15% by weight of a coating; wherein the core beads comprise about 18 to about 21% by weight of topiramate, about 8 to about 11% by weight of povidone, and about 58 to about 61% by weight of sugar spheres; and the coating comprises about 6 to about 9% by weight of cellulose acetate, and about 2 to about 5% by weight of povidone.

In a subclass of the invention is the pharmaceutical composition comprising about 89% by weight of core beads and about 11% by weight coating, wherein the core beads comprise about 19.8% by weight topiramate, about 9.9% by weight povidone, and about 59.3% by weight sugar spheres; and the coating comprises about 7.2% by weight cellulose acetate and about 3.8% by weight povidone.

Illustrative of the invention are methods of treating convulsions and/or epilepsy in a mammal in need thereof which comprises administering to the mammal a therapeutically effective amount of any of the pharmaceutical compositions of the present invention.

Also included in the invention are methods of treating a condition selected from neuropathic pain, amyotrophic lateral sclerosis, acute ischemia, obesity, diabetes, psoriasis or bipolar disorder (including manic depression) in a mammal in need thereof which comprises administering to the mammal a therapeutically effective amount of any of the pharmaceutical compositions of the present invention.

Detailed Description of the Invention

The present invention provides a solid dosage formulation of topiramate intended primarily for pediatric use, or for patients who cannot swallow tablets. More particularly, the solid dosage formulation is a sprinkle formulation comprising core particles of the active agent which is taste-masked with a second layer to obscure the extremely bitter taste of topiramate. The core particles can comprise

topiramate alone, e.g., in granular or crystalline form, or topiramate and one or more excipients which are then formed into granules or beads by techniques known to one of ordinary skill in the art, e.g., roller compaction and comminution, extrusion-spheronization or other methods of forming granules or beads. The
5 preferred solid dosage formulation of the present invention is in the form of microspheres which may be sprinkled onto soft food (e.g., baby food) and swallowed by the patient along with the food.

In a preferred embodiment, three strengths, 15, 25, and 50 mg, are
10 obtained from a single sprinkle formulation of topiramate coated onto sugar spheres using povidone as a binder, and taste-masked with a coating of cellulose acetate and povidone to form coated beads. The strengths are differentiated by means of differing fill weights and of proportional capsule sizes. That is, to aid in delivery of the appropriate dosage to the patient, an amount of coated beads
15 sufficient to deliver the desired dose may be encapsulated into a capsule, for example, a size 0, size 1, or size 2 gelatin capsule consisting of a white body with a natural cap. Black pharmaceutical ink can be utilized to provide product identification information on the capsules. For pediatric patients, the capsules can be opened and the contents of the capsules sprinkled onto food and ingested;
20 however, mature patients may swallow drug product in intact capsules, if desired.

In general, the process for the preparation of the sprinkle formulation includes a step in which core particles comprising granules, beads or crystals of topiramate, alone or in combination with one or more excipients, are coated with a
25 taste masking mixture and then dried. The term "particles" as used herein refers to free flowing substances of any shape which are larger than a powder including crystals, beads (smooth, round or spherical particles) and granules. A variety of methods known to those of ordinary skill in the art of pharmaceutical sciences may be employed to prepare the core particles comprising the active agent of
30 topiramate. In one method, granules or large single crystals of topiramate can be utilized as the core particles and coated with the taste masking mixture. The coated material formed from the granules or crystals of topiramate may then be

compressed into chewable tablets, if desired, or sprinkled onto soft food and swallowed.

In a second approach, the active agent of topiramate (in powder form) is first placed in a fluidized bed equipment and thereafter, a spray binder solution or suspension comprised of, for example, povidone, starch, sugar, syrup, HPMC among other excipients known to those of ordinary skill in the art in a pharmaceutically acceptable solvent (e.g., water, ethanol, acetone, among others) is sprayed onto the powder, formed into granules and then dried until the solvent is evaporated to provide the core particles. The drying temperature may vary over a broad range, but should not be so high as to render the active agent inactive. As a slight modification of this second approach, a suspension of topiramate and a binder in a pharmaceutically acceptable solvent is sprayed onto sugar spheres in a fluidized bed equipment and dried to provide core beads.

15

In a third method for forming the core particles, powdered or granular active agent, and diluent or bulking agent are mixed with water or a pharmaceutically acceptable solvent (e.g., water, ethanol) to form a wet mass. The mixture is mixed, e.g., in a Hobart mixer or other suitable mixer, until a wet mass or dough is formed. The wet mass is then placed in an extruder and extruded as a long thin strand. The mixture may then be dried and suitably comminuted or may be placed in a suitable spheronizer to make a pharmaceutical core that is round followed by drying. The drying temperature may vary over a broad range, but should not be so high as to render the active agent inactive.

20

Still another approach for forming the core particles is by roller compaction of topiramate, either alone or in combination with one or more excipients. For example, topiramate in powdered or granular form can be mixed with excipient to provide suitable binding and lubricity, for example microcrystalline cellulose, magnesium stearate or talc among others, and then passed through a compactor to compact the mixture into a mass. The mass is then passed through a size

25

reduction machine and reduced to a suitable particle size to provide the core particles.

As used herein, the term "topiramate" and "active agent of topiramate" are
5 synonymous and are used interchangeably throughout the specification to refer to the compound 2,3:4,5-bis-O-(1-methylethylidene)- β -fructopyranose sulfamate which forms the active agent of the pharmaceutical compositions of the present invention. Topiramate and its use for treating epilepsy and glaucoma are described in U.S. Patent No. 4,513,006. Topiramate can be synthesized
10 according to the processes disclosed in U.S. Patent Nos. 4,513,006 and 5,387,700, and preferably, according to the process of Examples 1-3 of U.S. Patent No. 5,387,700.

The term "therapeutically effective amount" as used herein means that
15 amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease being treated.

20 The term "excipient," as used herein, refers to any inert substance which may be combined with an active agent for preparing convenient dosage forms, including, for example, diluents, binders, lubricants, disintegrants, colors, flavors and sweeteners.

25 Suitable diluents for use in the formulation and processes of the present invention include, but are not limited to, dicalcium phosphate, calcium sulfate, lactose, sorbitol, microcrystalline cellulose, kaolin, mannitol, sodium chloride, dry starch, powdered sugar and sugar spheres. In a preferred embodiment of the invention, sugar spheres (20-60 mesh, preferably, 20-40 mesh, most preferably,
30 20-24 mesh) are utilized as a diluent in the core beads. In a particularly preferred embodiment, sugar spheres NF (20/25 mesh) available from Crompton & Knowles Corporation as NU-PAREIL PG® are used.

Suitable binders for use in the instant formulation and processes include, but are not limited to synthetic gums such as hydroxypropyl methylcellulose ("HPMC"), povidone, carboxymethylcellulose, ethylcellulose and methylcellulose, starch, pregelatinized starch, gelatin, sugars (e.g., molasses) and natural gums (e.g., acacia gum, sodium alginate, panwar gum). Preferably, povidone (especially, Povidone USP) is used as the binder. In a particularly preferred embodiment, the povidone is PLASDONE® (K29/32) supplied by ISP Technologies, Inc. as a manufacturer of GAF products.

10

Disintegrants which can be utilized in the formulation and processes of the present invention include, but are not limited to, methylcellulose, cellulose, carboxymethylcellulose, croscarmellose sodium, magnesium aluminum silicate, povidone, starch, sodium starch glycolate, pregelatinized starch, alginic acid and guar gum. Preferably, the disintegrant is povidone. In a particularly preferred embodiment, the povidone is PLASDONE® (K29/32) supplied by ISP Technologies, Inc. as a manufacturer of GAF products.

Suitable taste masking agents which can be used in the formulation and processes include, but are not limited to, cellulose acetate, cellulose acetate butyrate, ethylcellulose, methylcellulose (including ethylcellulose and methylcellulose combinations), and a wide range of copolymers available under the tradename of Eudragits (Röhm Pharma of Darmstadt, Germany). In a preferred embodiment, the taste masking agent is cellulose acetate (Cellulose Acetate, NF).

A variety of solvents may be used as the first and second solvent in the processes for preparing the pharmaceutical composition. Suitable solvents include, but are not limited to, water, acetone, alcohols (e.g., methanol, ethanol, isopropanol), methylene chloride, ethyl acetate, methyl ethyl ketone, and mixtures thereof. In a preferred embodiment, the first solvent used for forming the core beads is water, and the second solvent used for coating the core beads with the

taste masking mixture is an acetone-alcohol mixture, preferably, an acetone-ethanol mixture, more preferably, an acetone-dehydrated alcohol mixture.

In a preferred embodiment, a suspension of topiramate and a binder in a first solvent is sprayed onto sugar spheres (20-25 mesh) and dried to provide core beads. The core beads are then screened to remove fines and agglomerates. The core beads are coated again with a taste masking mixture and then dried. The taste masking mixture, which is sprayed onto the core beads, comprises a taste masking agent and a disintegrant dissolved or suspended in a second solvent, which may be the same or different from the first solvent. Coated beads are sifted to remove fines and agglomerates, prior to encapsulation.

In a particularly preferred embodiment of the process for preparing the sprinkle formulation, a suspension of topiramate in a solution of povidone in purified water is sprayed onto sugar spheres (20-25 mesh) and dried in a fluid bed processor equipped with a Wurster column. The ratio of topiramate:povidone utilized in the suspension can be 50:25, 50:30 or 50:35. Preferably, a 50:25 ratio of topiramate:povidone is used. The core beads then are screened to remove fines and agglomerates such that the core beads have a particle size between about 0.100 mm and about 2.5 mm, preferably, between about 0.5 mm and about 1.5 mm, most preferably, between about 0.710 mm and about 1.18 mm. The core beads are coated again with a taste masking mixture of cellulose acetate and povidone suspended in an acetone/alcohol mixture in a fluid bed unit equipped with a Wurster column, and dried. The ratio of cellulose acetate/povidone in the taste masking mixture can be 60/40, 50/50, 65/35 or 55/45; preferably, a 65/35 ratio of cellulose acetate/povidone is utilized. Coated beads are sifted to remove fines and agglomerates to provide a final particle size between about 0.100 mm and about 2.5 mm, preferably, between about 0.5 mm and about 1.5 mm, most preferably, between about 0.850 mm and about 1.18 mm. The coated beads are packaged (e.g., in capsules, sachets or other methods known to those of ordinary skill in the art) to deliver the desired amount of active ingredient to the patient.

When a particle size range is specified for the core and/or coated particles (e.g., between about 0.100 mm and about 2.5 mm), it is intended that at least 75%, preferably, 85%, and most preferably, 95% of the particles have a particle size falling within the specified range (e.g., about 0.100 mm and about 2.5 mm).

5

The invention will be described more specifically in terms of its preferred embodiment which is the preparation of a sprinkle formulation of topiramate. In the first step of the process, core beads are prepared by coating sugar spheres (20-25 mesh) with a suspension of topiramate and povidone in water. More particularly, the sugar spheres are placed in a fluidized bed coater and fluidized by a flow of warm air. The temperature of the air has not been found to be narrowly critical, and can vary over a wide range, however, the temperature should not be high enough to cause decomposition, sintering, or melting of the sugar spheres. When coating the sugar spheres with the topiramate/povidone suspension (preferably, a 50:25 ratio), a temperature from about 50° to 75°C has been found to be suitable. The rate of air flow is adjusted so as to fluidize the sugar spheres. Such flow will vary depending on factors such as the specific equipment used, the size of the individual sugar spheres, the size of the charge of sugar spheres, the apparent specific gravity of the spheres, and other factors known to the worker in the arts relating to fluidized bed coating. After the sugar spheres have been fluidized, a previously prepared suspension of topiramate in a solution of povidone in water is sprayed onto the fluidized bed to provide the core beads. The air flow through the bed is continued until the amount of water remaining in the topiramate core beads has been substantially reduced. The core beads are actually dry to the touch within a very short time after the topiramate suspension has been sprayed onto the sugar spheres. However, the total drying time required to ensure that the water content has been reduced to the desired level may take much longer, depending on the temperature of the air, the size of the batch, and the like. Routine experimentation will suffice to determine the appropriate air temperatures and total times required in the fluidized bed coaters in individual cases. The core beads are sized through a sifter using 16 mesh and 25 mesh screens.

In the second step of the process, the core beads are coated with a taste masking mixture to provide the coated beads of the sprinkle formulation. More specifically, the core beads are placed in a fluidized bed coater and fluidized by a flow of warm air. The temperature of the air has not been found to be narrowly critical, and can vary over a wide range, keeping in mind the fact that the temperature should not be so high as to cause decomposition, sintering, or melting, of the topiramate core beads. When coating the topiramate core beads, a temperature of from 30° to 75°C has been found to be suitable. The rate of air flow is adjusted so as to fluidize the core beads. Such flow will vary depending on factors such as the specific equipment used, the size of the charge of core beads, the size of the individual core beads, the apparent specific gravity of the core beads, and other factors known to the worker skilled in the arts of fluidized bed coating. After the core beads have been fluidized, a taste mask coating mixture is sprayed onto the fluidized bed. The taste mask coating mixture comprises a solution of cellulose acetate/povidone (preferably, in a 65:35 ratio) in an acetone-alcohol (preferably, acetone-dehydrated alcohol) solvent mixture. The air flow through the bed is continued until the amount of solvent remaining in the coating has been reduced to part per million levels. The coated beads are actually dried to the touch within a very short time after the coating solution has been sprayed onto the topiramate core beads. However, the total drying time required to ensure that the solvent content of the coating has been reduced to the level desired may take much longer, depending on the temperature of the air, the size of the batch and the like. Routine experimentation will suffice to determine the appropriate air temperature and total times required in the fluidized bed coaters in individual cases. The coated beads are then sized through a sifter using 16 mesh and 20 mesh screens.

A sprinkle formulation having satisfactory taste masking and bioavailability properties was obtained when the taste mask coating comprises about 7 to about 15% by weight of the final pharmaceutical composition. Preferably the taste mask coating comprises about 9% by weight to about 13% by weight, most preferably, about 11% by weight, of the pharmaceutical composition when dried.

Dissolution results in water indicative of bioavailability for the pharmaceutical composition having between 7 and 15% by weight taste mask coating are shown below in Table 1.

5

TABLE 1
DISSOLUTION RESULTS IN WATER

Coating (%)	10 min	20 min	30 min	45 min	60 min
(%) Dissolved					
7	35.0	72.9	91.2	98.4	99.2
9	26.8	58.1	84.3	97.7	100.8
11	21.7	52.3	79.1	97.3	99.7
13	15.5	40.9	66.3	91.4	98.8
15	12.5	35.3	59.7	85.6	96.6

10

To aid in delivery of the appropriate dosage to the patient, an encapsulation machine can be utilized to encapsulate an amount of coated beads to provide 15 mg, 25 mg and 50 mg strengths of topiramate into a size 2, 1 or 0 gelatin capsule, respectively.

15

While the use of fluidized bed coating has been described in some detail as one preferred method for making the core beads and the coated beads, other techniques for making the core and coated beads readily known to those of ordinary skill in the art may be used. Such other techniques include various microencapsulation techniques such as coacervation and solvent evaporation.

20

In a particularly preferred embodiment, the ingredients and amounts of each ingredient used to prepare the topiramate sprinkle bead formulation are provided in Table 2.

TABLE 2
TARGET COMPONENT/COMPOSITION

<u>INGREDIEN</u> <u>I</u>	<u>REFERENC</u> <u>E</u>	<u>ROLE</u>	<u>Unit Dosage Strength (mg)</u>		
			<u>50 mg</u>	<u>25 mg</u>	<u>15 mg</u>
Topiramate		Active	50.0	25.0	15.0
Povidone	USP	Binding Agent	25.0	12.5	7.5
Purified Water ¹	USP	Process Aide	—	—	—
Sugar Spheres, 20-25 mesh	NF	Core Bead	150.0	75.0	45.0
Cellulose Acetate	NF	Film Coat	18.076	9.038	5.423
Povidone	USP	Film Coat	9.733	4.8665	2.9199
Acetone ¹	NF	Process Aide	—	—	—
Dehydrated Alcohol ¹	USP	Process Aide	—	—	—
Gelatin Capsules	Type IV	Drug Product Holder	one unit (Size 0)	one unit (Size 1)	one unit (size 2)
Printing Ink		Identifier			

5 1. Essentially removed during drying.

The strengths of topiramate sprinkle capsules, 15, 25 and 50 mg are obtained from a single formulation of topiramate-coated beads by encapsulating the proportionate amounts of coated beads in appropriately sized and marked capsules. Table 3 provides a batch formula for a production batch of topiramate sprinkle bead formulation.

10

TABLE 3
BATCH FORMULATION

TARGETED FORMULATION

5

<u>INGREDIENT</u>	<u>TARGET (kg)</u>	<u>RANGE(kg)</u>	<u>RANGE (%)</u>
<u>CORE BEADS</u>			
Topiramate	37.5	—	—
Povidone, USP	18.75	±0.09375	±0.5%
Purified Water, USP ¹	93.75	±0.9375	±1.0%
Sugar Sphere, NF 20-25 mesh	112.50	—	—
Core Bead Batch Size	168.75		
<u>COATED BEADS</u>			
Core Beads	150.00		
Cellulose Acetate, NF	12.051	±0.12051	±1.0%
Povidone, USP	6.489	±0.06489	±1.0%
Acetone, NF ¹	120.00	±1.2%	±1.0%
Dehydrated Alcohol, USP ¹	30.00	±0.3%	±1.0%
Coated Bead Batch Size	168.54		
<u>GELATIN CAPSULES</u>			
Gelatin			
Print Ink			

1. Essentially removed during drying.

10 A comparison of dissolution rates in water between TOPAMAX® 100 mg tablets and topiramate sprinkle capsule formulations, 25 and 50 mg dosages (according to the specifications of Table 2) are shown in Table 4.

TABLE 4
DISSOLUTION COMPARISON

5

% Label Dissolved (mean)				
Product	10 min	20 min	30 min	45 min
100 mg Tablet	85.0 (79-89)	92.6 (89-96)	96.4 (93-99)	—
25 mg Sprinkle	19.7 (17-22)	51.4 (48-55)	75.0 (71-80)	94.7 (90-99)
50 mg Sprinkle	17.8 (17-19)	48.1 (45-50)	71.3 (69-73)	93.5 (91-96)

The stability of the sprinkle formulation of the present invention was compared to TOPAMAX® (topiramate) tablets by storing both formulations in controlled stability chambers for the purpose of determining the stability profile for the two products. Samples were stored at 30°C. The sprinkles were stored at 60% relative humidity (RH); relative humidity for the tablet batches was either controlled at 35% RH or was not controlled, but, in any event, was well below 60% RH. Data were collected for assay (amount of drug remaining), sulfate and sulfamate, physical appearance at selected time intervals, e.g., 18 months, 24 months. Physical appearance, i.e., discoloration to brown or black, and amount of sulfate detected are good indicators of degradation of the active agent (topiramate). For each mole of topiramate that degrades, a molar equivalent of inorganic impurity (sulfate/sulfamate) is formed. The presence of inorganic impurity is readily determined by one of ordinary skill in the art using standard techniques, e.g., HPLC.

At 18 months, some instability was detected by appearance data for the tablets, while the sprinkle formulation showed no signs of instability/degradation.

Clear signs of degradation were apparent by appearance and sulfate data for the tablets at 24 months. After 24 months of storage at 30 degrees 60% RH, the 25 and 50 mg strengths of sprinkle capsules remained stable while the 15 mg strength showed instability. At 25 degrees 60% RH storage for 24 months, all
5 three strengths of the sprinkle formulation remained stable.

It is known that moisture accelerates the degradation of topiramate. It has now been unexpectedly found that the coating used to taste mask the topiramate core beads also provides a barrier to the absorption of moisture, and therefore,
10 improves on the stability of the sprinkle formulation. For storage of the tablets, it was necessary to put a desiccant into the bottles to stabilize the tablet formulation. However, there is no need for a dessicant for the sprinkle formulation. In addition, the capsules which are used to ease delivery of the appropriate dosage of sprinkles contain more than 10% moisture by weight, and yet, this
15 moisture does not accelerate the degradation of topiramate because of the taste mask coating for the sprinkles.

The following examples are provided to further define the invention without, however, limiting the invention to the particulars of these examples.

20

Example 1

Preparation of Core Beads

25	<u>Ingredient</u>	<u>Batch Amount (kg)</u>
	Topiramate	37.50
	Povidone, USP	18.75
	Sugar Spheres, NF (20-24 mesh)	112.50
30	Purified Water, USP	93.70

Batch amounts of each of the core bead ingredients were accurately weighed out. In a jacketed kettle (approximately 60 gallons) equipped with a sweeper, a homogenizer (Silverson or equivalent) and a mixer (LIGHTNIN'® or equivalent) was placed the appropriate batch amount of purified water, USP. The batch amount of Povidone, USP was added and the resulting mixture mixed for a minimum of 15 minutes to disperse the povidone in the purified water. The topiramate (37.50kg) was added and the mixture mixed for a minimum of 15 minutes to disperse. Water was passed through the jacket. Using the mixer and homogenizer, the topiramate suspension was homogenized for approximately 90 minutes (range: 80-100 minutes). Stirring was continued through the steps which follow for preparing the core beads.

A pump (Masterflex or equivalent) was prepared with three pump heads for spraying. The batch quantity of sugar spheres, NF was charged to a fluid bed (Glatt Fluid Bed equipped with a 32 inch Wurster column, 3 guns with 2.2 mm nozzles, or equivalent). The sugar spheres were fluidized and the topiramate suspension sprayed through the nozzles (approximate spray rate: 1kg/min; approximate spray time: 2.25 hours) according to the parameters shown in Table 5.

20

Table 5

<u>Operating Function</u>	<u>Operating Parameter</u>
Air Flow	2400 CFM (range:1900-2900 CFM)
Inlet Air Temperature	60°C (range:50°C-70°C)
Bed Temperature	40°C (range:38°C-45°C)
Atomization Air	3 bar (range:2.7-3.5 bar)

The core beads were dried at 60°C for at least 15 minutes (range: 15-18 minute) after the bed temperature had reached 60°C (range: 55°C-65°C) according to the parameters provided in Table 6.

Table 6

Operating Function	Operating Parameter
Air Flow	2100 CFM (range: 1800-2200 CFM)
5 Inlet Air Temperature	60°C (range: 50°C-70°C)
Atomization Air	1 bar (range: 1-2 bar)

The core beads were then sized through a 48" sifter (Sweco or equivalent), using 16 mesh and 25 mesh screens to remove fines and agglomerates.

10

Example 2Preparation of Coated Beads

<u>Ingredient</u>	<u>Batch Amount (kg)</u>
15 Topiramate - Core Beads	150.00
Cellulose Acetate, NF	12.051
Povidone, USP	6.489
Acetone, NF	120.00
20 Dehydrated Alcohol, USP	30.00

The batch quantities of Acetone, NF and Dehydrated Alcohol, USP were transferred to a suitable stainless steel tank and mixed. The batch quantity of Povidone, USP was added using a suitable mixer (LIGHTNIN'® or equivalent).

25 The batch quantity of Cellulose Acetate, NF was added at the vortex while mixing with a suitable mixer and the coating solution was checked visually for clarity.

The 16/25 mesh topiramate core beads (150 kg) from Example 1 were fluidized in a Glatt Fluid Bed processor equipped with a Wurster column (or
30 equivalent). The core beads were sprayed with the coating solution until the entire quantity of coating solution was exhausted. The coated beads were dried at

approximately 60°C for a minimum of 30 minutes (range: 28-32 minutes).

Operating parameters are shown in Table 7.

Table 7

5

Operating Ranges for Spraying/Drying

<u>Parameter</u>	<u>Operating Range</u>
10 Inlet Air Flow	1500-3000 CFM
Inlet Air Temperature	30°C - 70°C
Atomization Air	1 - 4 bars
Bed Temperature	30°C - 70°C

- 15 A Sweco Sifter (or equivalent equipment) was fitted at the top with a 16 mesh screen and at the bottom with a 20 mesh screen. The entire batch of coated beads was sieved and coated beads outside the 16-20 mesh range discarded.

Example 3

20

Encapsulation of Coated Beads

An encapsulation machine (H&K Encapsulator or equivalent) was prepared with a bead filler attachment and the coated beads from Example 2 encapsulated.

- 25 Target fill weights were determined by assay of the coated beads prior to encapsulation. Variability of the fill weight was controlled by weight sorting, which is required for the 15 mg strength, but optional and employed as necessary for the 25 and 50 mg strengths. A KKE sorting machine (or equivalent) was used to weigh the filled capsules when weight sorting was employed. Filled capsules not
30 meeting the acceptable weight range were discarded by the sorter.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the

5 following claims and their equivalents.

WHAT IS CLAIMED IS:

1. A process for forming a pharmaceutical composition comprising:
 - (a) preparing core particles comprising an active agent of topiramate;
 - 5 (b) drying the core particles from step (a) to form dried core particles;
 - (c) coating the dried core particles from step (b) with a taste masking mixture to form coated particles; and
 - (d) drying the coated particles from step (c) to form the pharmaceutical composition wherein the amount of taste masking mixture ranges from about 7%
10 by weight to about 15% by weight of the pharmaceutical composition.
2. The process of Claim 1, wherein the core particles comprise the active agent of topiramate and at least one excipient.
- 15 3. The process of Claim 2, wherein the core particles comprise the active agent of topiramate, a binder and a diluent wherein the diluent is sugar spheres.
4. The process of Claim 3, wherein the taste masking mixture
20 comprises about 9 to about 13% by weight of the pharmaceutical composition.
5. The process of Claim 4, wherein the dried core particles of step (b) are sized to between about 0.100 mm and about 2.5 mm prior to coating with the taste masking mixture.
25
6. The process of Claim 5, wherein the coated particles from step (d) are sized to between about 0.100 mm and about 2.5 mm.
7. The process of Claim 6, wherein the core particles are prepared by
30 spraying a suspension of topiramate and the binder in a solvent onto the sugar spheres.

8. The process of Claim 7, wherein the binder is selected from povidone, HPMC, acacia gum, sugar, molasses, sodium alginate, panwar gum, starch, pregelatinized starch, carboxymethylcellulose, ethylcellulose or methylcellulose.

5

9. The process of Claim 8, wherein the binder is povidone.

10. The process of Claim 8, wherein the taste masking mixture comprises a taste masking agent selected from cellulose acetate, cellulose acetate buytrate, methylcellulose, ethylcellulose or a Eudragit; and a disintegrant selected from povidone, methylcellulose, starch, sodium starch glycolate, pregelatinized starch, cellulose, carboxymethylcellulose, croscarmellose sodium, magnesium aluminate silicate, alginic acid or guar gum.

15

11. The process of Claim 10, wherein the taste masking mixture comprises cellulose acetate and povidone.

12. The process of Claim 11, further comprising encapsulating the coated bead.

20

13. A pharmaceutical composition made by the process of Claim 1.

14. A pharmaceutical composition comprising

25

(a) core particles containing an active agent of topiramate, wherein the core particles have an initial particle size between about 0.100 mm and 2.5 mm; and

(b) a taste mask coating, wherein the taste mask coating comprises between about 7% by weight and about 15% by weight of the pharmaceutical composition and wherein the coated particles of the pharmaceutical composition have a final particle size of about 0.100 mm to about 2.5 mm.

30

15. The pharmaceutical composition of Claim 14, wherein the core particles comprise the active agent of topiramate and at least one excipient.

16. The pharmaceutical composition of Claim 15, wherein the core particles comprise the active agent of topiramate, a binder and a diluent wherein the diluent is sugar spheres.

17. The pharmaceutical composition of Claim 16, wherein the taste mask coating comprises between about 9% by weight and about 13% by weight of the pharmaceutical composition.

18. The pharmaceutical composition of Claim 17, wherein the taste mask coating comprises about 11% by weight of the pharmaceutical composition.

19. The pharmaceutical composition of Claim 18, wherein the core particles have an initial particle size between about 0.5 mm and 1.5 mm and the coated particles of the pharmaceutical composition have a final particle size between about 0.5 mm and 1.5 mm.

20. The pharmaceutical composition of Claim 19, wherein the core particles have an initial particle size between about 0.710 mm and 1.18 mm and the coated particles of the pharmaceutical composition have a final particle size between about 0.850 mm and 1.18 mm.

21. The pharmaceutical composition of Claim 20, wherein the binder is selected from povidone, HPMC, sodium alginate, panwar gum, acacia gum, gelatin, sugar, molasses, starch, pregelatinized starch, methycellulose, ethylcellulose or carboxymethylcellulose; and the taste mask coating comprises a taste masking agent and a disintegrant, wherein the taste masking agent is selected from cellulose acetate, methylcellulose, ethylcellulose, a Eudragit or cellulose acetate butyrate; and the disintegrant is selected from povidone, cellulose, carboxymethylcellulose, croscarmellose sodium, magnesium aluminate

silicate, starch, sodium starch glycolate, pregelatinized starch, alginic acid or guar gum.

22. The pharmaceutical composition of Claim 21, wherein the binder is
5 povidone, the taste masking agent is cellulose acetate and the disintegrant is povidone.

23. The pharmaceutical composition of Claim 22, wherein the coated
particles of the pharmaceutical composition are encapsulated.

10

24. A pharmaceutical composition comprising about 85 to about 93% by
weight core beads, and about 7 to about 15% by weight of a coating; wherein the
core beads comprise about 18 to about 21% by weight of topiramate, about 8 to
about 11% by weight of povidone, and about 58 to about 61% by weight of sugar
15 spheres; and the coating comprises about 6 to about 9% by weight of cellulose
acetate, and about 2 to about 5% by weight of povidone.

25. The pharmaceutical composition of Claim 24, comprising about 89%
by weight of core beads and about 11% by weight coating, wherein the core beads
20 comprise about 19.8% by weight topiramate, about 9.9% by weight povidone, and
about 59.3% by weight sugar spheres; and the coating comprises about 7.2% by
weight cellulose acetate and about 3.8% by weight povidone.

26. A method of treating convulsions in a mammal in need thereof which
25 comprises administering to the mammal a therapeutically effective amount of the
pharmaceutical composition of Claims 14.

27. A method of treating epilepsy in a mammal in need thereof which
comprises administering to the mammal a therapeutically effective amount of the
30 pharmaceutical composition of Claims 14.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 September 1999 (10.09.1999)

PCT

(10) International Publication Number
WO 99/044581 A3

- (51) International Patent Classification⁶: **A61K 9/16**, 9/28, 31/35, 31/70, 31/095
- (21) International Application Number: **PCT/US99/04449**
- (22) International Filing Date: **1 March 1999 (01.03.1999)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/076,770 **4 March 1998 (04.03.1998)** **US**
- (71) Applicant (for all designated States except US): **OR-THO-McNEIL PHARMACEUTICAL, INC.** [US/US]; U.S. Route #202, Raritan, NJ 08869-0602 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **THAKUR, Madhav, S.** [US/US]; 112 Mendham Drive, North Wales, PA 19477 (US). **KOTWAL, Pramod, M.** [US/US]; 145 Cambronne Circle, Blue Bell, PA 19422 (US). **GIBBS, Irwin, S.** [US/US]; 144 Roy Lane, Huntington Valley, PA 19006 (US).
- (54) Title: **PHARMACEUTICAL COMPOSITION OF TOPIRAMATE**
- (57) Abstract: The invention is directed to a pharmaceutical composition of topiramate, an anticonvulsant which is useful for treating epilepsy. More specifically, the present invention provides a solid dosage formulation of topiramate intended primarily for use by pediatric patients, of for patients who have difficulty swallowing tablets. Processes for preparing the pharmaceutical composition are also described.
- (81) Designated States (national): **AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW.**
- (84) Designated States (regional): **ARIPO** patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), **Eurasian** patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), **European** patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), **OAPI** patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
17 July 2003
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 99/044581 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/04449

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K9/16 A61K9/28 A61K31/35 A61K31/70 A61K31/095

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 88 03795 A (MEHTA ATUL M) 2 June 1988 (1988-06-02) page 1, line 5 - line 17 page 7 -page 20 ---	1-25
A	EP 0 350 701 A (RESA FARMA) 17 January 1990 (1990-01-17) page 2, line 42 - line 45 page 3, line 10 - line 56 ---	1-25
A	EP 0 138 441 A (MCNEILAB INC) 24 April 1985 (1985-04-24) page 7, line 33 -page 8, line 26 abstract -----	1,13

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 June 1999

Date of mailing of the international search report

6. 07. 99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mosser

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/04449

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **26 and 27**
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 26 and 27 concern a method for treatment of the human or animal body by therapy (see rule 39.1 (iv) PCT), the search has been carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/04449

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 8803795	A	02-06-1988	US	4800087 A		24-01-1989
			AT	75605 T		15-05-1992
			DE	3778892 D1		11-06-1992
			DK	406788 A		16-09-1988
			EP	0302900 A1		15-02-1989
			JP	1502589 T		07-09-1989
			WO	8803795 A1		02-06-1988

EP 0350701	A	17-01-1990	IT	1226549 B		24-01-1991
			EP	0350701 A2		17-01-1990
			JP	2059515 A		28-02-1990

EP 0138441	A	24-04-1985	US	4513006 A		23-04-1985
			AT	36149 T		15-08-1988
			AU	564842 B2		27-08-1987
			AU	3350484 A		04-04-1985
			CA	1241951 A1		13-09-1988
			DE	3473143 D1		08-09-1988
			DK	198191 A ,B,		09-12-1991
			DK	198291 A		09-12-1991
			DK	457784 A ,B,		27-03-1985
			EP	0138441 A2		24-04-1985
			ES	8602634 A1		16-03-1986
			FI	843765 A ,B,		27-03-1985
			HU	36784 A2		28-10-1985
			IE	57684 B1		24-02-1993
			JP	1804249 C		26-11-1993
			JP	5005824 B		25-01-1993
			JP	60109558 A		15-06-1985
			JP	5331132 A		14-12-1993
			KR	9201775 B1		02-03-1992
			MX	9202630 A1		30-06-1992
			NO	843836 A ,B,		27-03-1985
			NZ	209494 A		06-03-1987
			US	4582916 A		15-04-1986
			ZA	8407550 A		28-05-1986
